

The genus *Solidago* is characterized by a *naked receptacle, the down simple, rays of the corolla about five, scales of the calyx imbricated and close.* It is a very natural genus, easily distinguished at sight by its crowded tufts of compound flowers, which are almost always of a deep golden yellow.\*

The species *odora* has its *stem nearly smooth, leaves linear-lanceolate, entire, smooth, with a rough margin, and covered with pellucid dots. Racemes paniced, one sided.*

Class *Syngenesia*,—Order *Superflua*,—Natural orders *Compositæ*, Lin. *Corymbiferae*, Juss.

The sweet scented Golden rod grows in woods and fields throughout the United States, and flowers in September. It has a smooth appearance, and is among the smaller species of its family. The root is woody, much branched and creeping. Stem slender, from two to three feet high, smooth or slightly pubescent below, pubescent at top. The leaves are linear-lanceolate, closely sessile, broad at base, entire, acute, with only the midrib distinct, rough at the margin but otherwise smooth, and covered with pellucid dots, like *Hypericum perforatum*. The flowers grow in a compound, paniced raceme, with each of its branches

\* The only exception which I now recollect is *Solidago bicolor*, whose ray is white.

supported by a small leaf. These branches or peduncles are very slender and rigid, each giving off a row of ascending, downy pedicels, with small, linear bractes at their bases. Scales of the calyx oblong, acute, smooth, or slightly pubescent, the lower ones shorter and closely imbricating the rest. Florets of the ray few, with oblong, obtuse, ligules. Those of the disc funnel shaped, with acute segments. Down simple to the naked eye, feathery under the microscope. Seeds oblong.

This plant is the *Solidago odora* of Muhlenberg, and agrees with the character of Aiton. The *Solidago odora* of Michaux is possibly a different species. Willdenow's plant was undoubtedly different. The *folia puncticulosa*, which constitutes so distinct a mark in this species, I have not seen noticed by any botanist.

The leaves of the *Solidago odora* have a delightfully fragrant odour, partaking of that of anise and sassafras, but different from either. When subjected to distillation, a volatile oil, possessing the taste and aroma of the plant in a high degree, collects in the receiver. This oil apparently has its residence in the transparent cells, which constitute the dotting of the leaves, for the root is wholly destitute of the peculiar fragrance of the herb, and has rather a nauseous taste. This is contra-